EPJ manuscript No.
(will be inserted by the editor)

arXiv:1210.1323v1 [hep-ph] 4 Oct 2012

Wounded Quarks and Multiplicity at Relativistic Ion Colliders

Ashwini Kumar[1a], B. K. Singh[1], P. K. Srivastava[1b], and C. P. Singh[1]

Department of Physics, Banaras Hindu University, Varanasi 221005, INDIA

Received: date / Revised version: date

Abstract. In this paper, we propose a parameterization which is based on a phenomenological model involving the wounded quarks interactions for explaining the average charged particle multiplicity $\langle n_{ch}\rangle$ and the central pseudo-rapidity density $\langle (dn/d\eta)_{\eta=0}\rangle$ and their variations with the center-of-mass energies ($\sqrt{s_{NN}}$) in relativistic heavy-ion collider experiments. The model also interrelates nucleus-nucleus (A-A) collisions with p-A and p-p interactions. Our parameterization rests on simple assumptions regarding mean number of participating quarks and their average number of collisions. The results for $\langle n_{ch}\rangle$ and their variations with the mass number of colliding nuclei and $\sqrt{s_{NN}}$ are well supported by the experimental data. The results obtained from modified Glauber model involving wounded nucleons are also shown for comparison. Finally, we give the predictions from our model for A-A collisions at the Large Hadron Collider (LHC) and Compressed Baryonic Matter (CBM) experiments. Our results indicate towards a possible universal production mechanism existent for p-p, p-A and A-A collisions.

PACS. 25.75.Dw – 25.75.Gz – 25.75.-q – 12.38.Mh

1 Introduction

The ultimate goal of ultra-relativistic heavy-ion collision experiments is to test the predictions of quantum chromodynamics (QCD) because these can reveal the nature of hadronic interactions at extreme temperature and/or density and throw light on the role played by the quarks and gluons in multiparticle production mechanism [1]. In order to verify the existence of a deconfined plasma created in nuclear collisions, we require accurate and precise knowledge about its subsequent hadronization. Hadronic multiplicities and their correlations can reveal enough information on the nature, composition and size of the fireball formed in the collider experiments [2,3]. The evolution of the measured global observables in such experiments varies with the size and mass of the colliding nuclei and it emphasizes that significant rescatterings or multiple scatterings occur and contribute to the evolution of the energy deposition in the fireball.

The search for some regularities and systematics present in the average multiplicity distributions of hadron-hadron (h-h), hadron-nucleus (h-A) and nucleus-nucleus (A-B) collisions has grown into a fascinating topic in strong interaction physics today because these properties hint at the underlying universal production mechanism for charged particles. The study of $p-p$ and $p-A$ collisions is particularly useful because it helps our understanding, e.g., in which way $A-B$ collisions differ from simple superposition of nucleon-nucleon collisions. However, our understanding of the subject is still quite poor partially because of the reason that in soft hadronic or nuclear collisions, the role of perturbative QCD is less clear as it fails close to the phase hadronization point where non-perturbative effects become clearly dominant. Several new experimental information on the multiparticle production have been reported in the recent past. Consequently a lot of efforts have also been put forward to understand and/or organise the experimental data by using various theoretical as well as phenomenological schemes [4,5,6,7,8,9,10, 11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26]. Some of these models explain multiplicities in nucleus-nucleus collisions mostly by using either quark-quark interactions or by considering basic nucleon-nucleon scatterings [23, 24,25,26]. However, predictions of models that successfully describe multiparticle production at RHIC energy usually differ at LHC energy by a factor of two [27] which signifies a major discrepency. So we focus our efforts in formulating a model which can describe multiplicity distributions consistently in the entire energy range accesible today at Relativistic Heavy Ion Collider (RHIC), Large Hadron Collider (LHC) and/or in future at Compressed Baryonic Matter (CBM) experiment of the FAIR (Facility of Antiproton and Ion Research) at GSI Darmstadt.

How one can describe the mechanics of such a relativistic many body problem in terms of simple, basic physical principles pose a major challenge in high energy heavy-ion physics today. Nucleus-nucleus collisions differ from

---

[a] *e-mail : ashwini.physics@gmail.com*
[b] *e-mail : prasu111@gmail.com*

nucleon-nucleon collisions in many respects. The presence of many constituent quarks participating in a chain of collisions makes the picture too complicated. In this paper, we describe nucleus-nucleus collisions in terms of effective number of constituent quarks participating in the collisions and the effective number of collisions suffered by each of them. In this model, we propose a gluon being exchanged between a quark of projectile or first nucleus and a quark belonging to target or other colliding nucleus. The resulting colour force is thus somewhat stretched between them and other constituent quarks because they try to restore the colour singlet behaviour. When two quarks separate, the colour force builds up a field between them and as the energy in the colour field increases, the colour tubes thus formed finally break-up into new hadrons and/or quark-antiquark pairs. We have essentially modified the phenomenological model earlier proposed by Singh et al [28]. We consider a multiple collision scheme in which a valence quark of the incident nucleon suffers one or more inelastic collisions with the quarks of target nucleons. In this process, quark loses energy and momentum which results into creation of new hadrons. Further, we assume that all the quark collisions are independent and their effects are incoherently superimposed. We find that charged particle multiplicities arising from $p-p$, $p-A$ and $A-B$ collisions can be accounted in a consistent manner by adequately incorporating the number of participating quarks and mean number of quark-quark collisions. The rest of the paper is organised as follows. In section II, the detailed description of the model is presented, the results and discussions are summarized in section III and finally in section IV, main conclusions and future prospects of this study are succinctly given.

2 Description of Model

If we assume a universal mechanism of charged particle production in the hadron-hadron, hadron-nucleus and nucleus-nucleus collisions, it must be driven by the available amount of energy involved for the secondary production, and it must depend on the mean number of participant quarks [29]. The main ingredients of our model are taken from the paper by Singh et. al. [28]. There exist several papers in the literature emphasizing various types of fitting parametrizations in order to provide a unified description for the produced charged particles in $p-p$ collisions at various energies. We have taken some of these functions with the values of their constants from the literature [30,31,32,33] and fitted them with the experimental data [34,35,36,37,38,39,40,41,42] as shown in Fig. 1. All these parametrizations fit the experimental data for $p-p$ collisions at intermediate and higher energies but usually show some disagreement with the experimantal data at lower energies (see Fig. 1). Recently, Jeon and collaborators have shown that the total mean multiplicity obtained at RHIC can be bounded by a cubic logarithmic term in the energy [43]. Based on the findings of Ref. [43], we propose here a new parameterization which extends upto a cubic logarithmic term to accomodate the entire $p-p$ experimental data [34,35,36,37,38,39,40,41,42] starting from low energies (i.e. from 6.15 GeV) upto the highest LHC energy (i.e. 7 TeV) as follows :

$$< n_{ch} >_{hp}= (a'+b'ln\sqrt{s_a}+c'ln^2\sqrt{s_a}+d'ln^3\sqrt{s_a})-\alpha. \quad (1)$$

In Eq. (1), α is the leading particle effect and $\sqrt{s_a}$ is the available center-of-mass energy (i.e., $\sqrt{s_a} = \sqrt{s} - m_B - m_T$ [33], where m_B is the mass of projectile and m_T the mass of the target nucleon, respectively), a', b', c' and d' are chosen constants. We take the values $a' = 1.8$, $b' = 0.37$, $c' = 0.43$ and $d' = 0.04$ as derived from the best fit to the data. The value of α is taken as 0.85 in our calculations [28].

We can extrapolate the validity of this parametrization for the produced charged particles in the hadron-nucleus interaction by considering the multiple collision effect in the nucleus. The number of constituent quarks which participated in hadron-nucleus (h-A) collisions share the total available center-of-mass energy $\sqrt{s_{hA}}$ and thus the energy available for each interacting quark becomes $\sqrt{s_{hA}}/N_q^{hA}$ [28], where N_q^{hA} is the mean number of constituent quarks in h-A collisions. The total available squared center-of-mass energy s_{hA} in h-A collisions can be related to s_a as $s_{hA} = \nu_q^{hA} s_a$ [44,45] with ν_q^{hA} as the mean number of inelastic collisions of quarks with target nucleus of atomic mass A. Within the framework of the Additive Quark Model [46,47,48,49,50], the mean number of collisions in hadron-nucleus interactions is discussed in relation to the ability of constituent quarks in the projectile hadron to interact repeatedly inside a nucleus. The average number of constituent quarks (N_q^{hA}) interacting inelastically in h-A collisions is given by [51],

$$N_q^{hA} = \frac{\bar{\nu}(A)}{\bar{\nu}_{qA}}, \quad (2)$$

where, $\bar{\nu}(A)$ is the mean number of collisions of the incident hadron and is equivalent to [51],

$$\bar{\nu}(A) = \bar{\nu}_{qA} \frac{N_c \sigma_{qA}^{in}}{\sigma_{hA}^{in}}, \quad (3)$$

with σ_{hA}^{in} as the inelastic hadron-nucleus cross-section and N_c is the number of valence quarks, $\bar{\nu}_{qA}$ represents the mean number of collisions of the wounded quark inside the nucleus A and is defined by [51],

$$\bar{\nu}_{qA} = \frac{A\sigma_{qN}^{in}}{\sigma_{qA}^{in}}, \quad (4)$$

where σ_{qN}^{in} and σ_{qA}^{in} are the inelastic cross-sections for quark-nucleon (q-N) and quark-nucleus (q-A) interactions, respectively and A is the atomic mass of the target nucleus.

Now, using the Eq. (2), (3) and (4), the average number of constituent quarks (N_q^{hA}) in h-A collisions can be rewritten as [28,51],

$$N_q^{hA} = \frac{N_c \sigma_{qA}^{in}}{\sigma_{hA}^{in}}. \quad (5)$$

The quark-nucleus inelastic cross-section (σ_{qA}^{in}) is determined from σ_{qN}^{in} ($\approx \frac{1}{3}\sigma_{NN}^{in}$) by using Glauber's approximation given as follows [28]:

$$\sigma_{qA}^{in} = \int d^2b \left[1 - \left(1 - \sigma_{qN}^{in} D_A(b)\right)^A\right], \tag{6}$$

where profile function $D_A(b)$ is related to the nuclear density, $\rho(b,z)$ by the relation [28] :

$$D_A(b) = \int_{-\infty}^{\infty} \rho(b,z)dz, \tag{7}$$

and we have chosen the form for the nuclear density as follows [52]:

$$\rho(b,z) = \frac{\rho_0}{1 + exp(\frac{\sqrt{b^2+z^2}-R}{a})}, \tag{8}$$

where all the notations have their usual meaning [52]. We have taken the values of the constants i.e., ρ_0, R and a for different colliding nuclei separately from the Ref. [53].

By using Eq. (5), one can easily see that the average number of constituent quark (N_q) is 1 for h-p collisions as in nucleon-nucleon collisions the interactions occur between single constituent or dressed quark in accordance with the additive quark picture [49,50] and other quarks are considered to be spectators. Thus the energy of the initial thermalized state which is responsible for the production of secondary particles is mainly due to the interacting single quark pair. The spectator quarks which are not the part of thermalized volume at the moment of collision, do not participate in secondary particle production process. As a result, the leading particles [54,55] resulting from the spectator quarks carry away a significant part of the energy [29]. Finally, the expression for average charged hadron multiplicity in h-A collisions can be expressed as follows [28]:

$$< n_{ch} >_{hA} = N_q^{hA} \left[a' + b'ln\left(\frac{\sqrt{s_{hA}}}{N_q^{hA}}\right) + c'ln^2\left(\frac{\sqrt{s_{hA}}}{N_q^{hA}}\right) + d'ln^3\left(\frac{\sqrt{s_{hA}}}{N_q^{hA}}\right)\right] - \alpha. \tag{9}$$

The generalization of the above picture for the case of nucleus-nucleus collisions can be achieved as follows [28]:

$$< n_{ch} >_{AB} = N_q^{AB} \left[a' + b'ln\left(\frac{\sqrt{s_{AB}}}{N_q^{AB}}\right) + c'ln^2\left(\frac{\sqrt{s_{AB}}}{N_q^{AB}}\right) + d'ln^3\left(\frac{\sqrt{s_{AB}}}{N_q^{AB}}\right)\right], \tag{10}$$

where $\sqrt{s_{AB}} = A\sqrt{\nu_q^{AB} s_a}$ [28] and the mean number of inelastic quark collisions ν_q^{AB} is [28]:

$$\nu_q^{AB} = \nu_{qA}\nu_{qB} = \frac{A\sigma_{qN}^{in}}{\sigma_{qA}^{in}}.\frac{B\sigma_{qN}^{in}}{\sigma_{qB}^{in}}. \tag{11}$$

Furthermore, mean number of participating quarks N_q^{AB} in A-B collisions can be calculated by generalizing Eq. (8) in the following manner [28]:

$$N_q^{AB} = \frac{1}{2}\left[\frac{N_B\sigma_{qA}^{in}}{\sigma_{AB}^{in}} + \frac{N_A\sigma_{qB}^{in}}{\sigma_{AB}^{in}}\right], \tag{12}$$

where σ_{AB}^{in} is the inelastic cross-section for nucleus-nucleus (A-B) collision and can be expressed in the following manner [28,56]:

$$\sigma_{AB}^{in} = \pi r^2 \left[A^{1/3} + B^{1/3} - \frac{c}{A^{1/3} + B^{1/3}}\right]^2, \tag{13}$$

where c is a constant and has a value 4.45 for nucleus-nucleus collisions. One can see that the extended parametrization in Eq. (10) relates nucleus-nucleus collisions to hadron-nucleus and hadron-proton collisions and the values of the parameters a', b', c', and d' remain unaltered which shows a universality in the role of basic quark-gluon interactions in all these processess together.

In a search for creating quark gluon plasma (QGP), greater emphasis is laid on the central or head-on collisions of two nuclei. The resulting mean multiplicity in central collisions can be obtained by using Eq. (10) in the following manner :

$$< n_{ch} >_{AB}^{central} = A \left[a' + b'ln(\nu_q^{AB} s_a)^{1/2} + c'ln^2(\nu_q^{AB} s_a)^{1/2} + d'ln^3(\nu_q^{AB} s_a)^{1/2}\right]. \tag{14}$$

The pseudorapidity distribution of charged particles is another very important quantity in the studies of particle production mechanism in high energy $h-h$ and $A-B$ collisions, which, however, is still not understood properly. It has been pointed out that $(dn_{ch}/d\eta)$ can be used to get the information on the temperature (T) and density (ρ) of the QGP [57,58,59]. Moreover, mid-rapidity density reflects the different stages of the reactions. To calculate the pseudorapidity density of charged hadrons, we first fit the experimental data of $(dn_{ch}/d\eta)_{\eta=0}^{pp}$ by using a similar parametrization as used earlier in Eq. (1), but with slightly different values of constants. Further, we calculate the central pseudo-rapidity density $(dn_{ch}/d\eta)_{\eta=0}^{AB}$ by using the basic idea of two component model for nuclear collsions [60], which suggests that the pseudo-rapidity density for $A-B$ collisions comes from two parts i.e., contributions arising due to hard processes and the other due to the soft processes. The contributions from hard processes is propotional to the total number of collisions i.e., $N_q^{AB}\nu_q^{AB}$ and the portion from soft processes is propotional to the participating quark constituents i.e., N_q^{AB} only. Thus, the expression for $(dn_{ch}/d\eta)_{\eta=0}^{AB}$ can be parameterized in terms of p-p rapidity density as follows :

$$\left(\frac{dn_{ch}}{d\eta}\right)_{\eta=0}^{AB} = \left(\frac{dn_{ch}}{d\eta}\right)_{\eta=0}^{pp} \left[(1-x)\, N_q^{AB} + xN_q^{AB}\nu_q^{AB}\right], \tag{15}$$



Fig. 1. Variations of total mean multiplicities of charged hadrons in $p-p$ collisions with $\sqrt{s_{NN}}$.



Fig. 3. Variations of total mean multiplicities of charged hadrons in central collisions for different colliding nuclei with $\sqrt{s_{NN}}$.



Fig. 2. Variations of total mean multiplicities of charged hadrons in central $Au-Au$ collisions with $\sqrt{s_{NN}}$. Experimental data points are taken from Ref. [62,63,64,65].

where x quantifies the relative contributions of hard and soft processes. The fraction x corresponds to the hard processes and the remaining fraction $(1-x)$ thus describes the contributions from the soft processes. In Eq. (15), we have used the parameterization for $(dn_{ch}/d\eta)^{pp}_{\eta=0}$ same as used in Eq. (1) for calculating $< n_{ch} >$ in p-p collisions and can be given as follows:

$$< (dn_{ch}/d\eta)^{pp}_{\eta=0} > = (a_1^{'} + b_1^{'} ln\sqrt{s_a} + c_1^{'} ln^2\sqrt{s_a} + d_1^{'} ln^3\sqrt{s_a}) - \alpha, \tag{16}$$

where $a_1' = 1.78$, $b_1' = 0.025$, $c_1' = 0.041$ and $d_1' = 0.0017$ as derived from the best fit to the data. We take the value of x at different $\sqrt{s_{NN}}$ according to Ref. [60].

3 Results and Discussions

In Fig. (1), we present the inelastic (filled symbols) and non-single diffractive (NSD) data (open symbols) of charged hadron multiplicity in full phase space for $p-p$ collisions at various center-of-mass energies from different experiments e.g., ISR, UA5 and E735 [34,35,36,37,38,39,40,41, 42]. We use inelastic data at very low energies (filled symbols) because NSD data are not available for these energies and also the trend shows that the difference between inelastc and NSD data is very small at lower energies. Further, we fit this data set with four different functional forms. The short-dashed line has the functional form as : $a_2 + b_2 s^{1/4}$ which is actually inspired by the Fermi-Landau model [30,31,32]. It provides a resonable fit to the data at higher $\sqrt{s_{NN}}$ with $a_2 = 5.774$ and $b_2 = 0.948$ [40]. However, since a_2 summarizes the leading particle effect, its value should not lie much larger than two. The dotted line represents the functional form as : $a_1 + b_1\ lns + c_1\ ln^2 s$ [33] and it fits the data well at higher $\sqrt{s_{NN}}$ but shows a large disagreement with the experimental data at lower center-of-mass energies. The dashed-dotted line represents the form $a_3 + b_3 s^n$ [33] and it also provides a good qualitative description of the data with $a_3 = 0$, $b_3 = 3.102$ and $n = 0.178$ [40]. The solid line represents our parametrization given by Eq. (1) and it undoubtedly offers the most reasonable fit to the data starting from very low energy upto very high $\sqrt{s_{NN}}$.

Fig. 2, shows the variation of mean multiplicity of charged hadrons produced in central $Au-Au$ collision with respect to $\sqrt{s_{NN}}$. We also compare our model results (solid line) and the modified Glauber model results (dash-dotted line) [61] with the experimental data of AGS and RHIC [62,63,64,65]. We would like to mention here the main difference between modified Glauber model and the standard Glauber calculation. In standard Glauber model, a participating nucleon can have one or more number of collision at the same $\sqrt{s_{NN}}$ and with the same value of $\sigma^{pp}_{inel}(\sqrt{s_{NN}})$. However, in modified Glauber model, a participating nucleon loses a fraction of its momentum after the first collision and participate in subsequent collisions with somewhat lower energy and also at a lower value of $\sigma^{pp}_{inel}(\sqrt{s})$. This modification significantly suppresses the

Table 1. Total mean multiplicity of charged hadrons produced in $d-Au$ collision at $\sqrt{s_{NN}}=200$ GeV.

	Colliding Nuclei	$\langle n_{ch}\rangle^{min.bias}$	$\langle n_{ch}\rangle^{central}$
Our Model	$d-Au$	82	162
PHOBOS [67]	$d-Au$	87^{+7}_{-6}	167^{+14}_{-11}



Fig. 4. Variations of pseudorapidity density at mid-rapidity for $p-p$ collisions with $\sqrt{s_{NN}}$. Filled symbols are data from inelastic $p-p$ events and open symbols are experimental data for NSD events [35,36,37,38,42,68,69,70,71,72]. Solid line is the outcome of our parametrization.

number of collisions (N_{coll}) in comparison to that obtained in the standard Glauber model. We find that the results obtained in our model give a good description to the experimental data in comparison to the modified Glauber model predictions, especially at lower center-of-mass energies.

In Fig. 3, we show the variation of mean multiplicity calculated from our model for $Au-Au$, $Pb-Pb$ and $Cu-Cu$ central collisions with respect to $\sqrt{s_{NN}}$. We also compare our model results with the experimantal data of RHIC and SPS experiments [62,63,64,65,66]. One can see from Fig. 3, that the model results give excellent fit to the separate experimental data points for $Cu-Cu$, $Au-Au$ and $Pb-Pb$, respectively. This concludes that our phenomenological model works well because it considers the picture of incoherent superposition of basic quark-quark collisions in describing the nucleus-nucleus collisions.

In Table I, we have shown the mean multiplicity of charged hadrons in minimum-bias as well as in central $d-Au$ collisions at $\sqrt{s_{NN}}=200$ GeV as obtained in our model. We have also shown the experimental results [67] for this collision for comparison. The good agreement between the experimental and model results shows that it can be used for asymmetric colliding nuclei as well.

In Fig. 4, we present the inelastic (filled symbols) and non-single diffractive (NSD) data (open symbols) of $\frac{dn_{ch}}{d\eta}$ at mid-rapidity for $p-p$ collisions at various center of mass energies from different experiments e.g., ISR, UA5, E735, RHIC and LHC [35,36,37,38,42,68,69,70,71,72]. Further, we fit (solid line) this data set with our functional form of Eq. (16), which is similar to Eq. (1) and obtained the values of constants as $a_1^{'}=1.78$, $b_1^{'}=0.025$, $c_1^{'}=0.041$ and $d_1^{'}=0.0017$. The central rapidity hadron density thus continues to increase with the colliding energy at LHC by an added triple logarithmic form.



Fig. 5. Variations of pseudorapidity density at mid-rapidity for $A-A$ collision with $\sqrt{s_{NN}}$. Symbols are data from various heavy ion collision experiments [62,63,64,65,66,73,74].

Table 2. Predictions of total mean multiplicity of charged hadrons produced in heavy-ion collision experiments.

Experiment	LHC	CBM I	CBM II
Colliding Nuclei	$Pb-Pb$	$Au-Au$	$Au-Au$
$\sqrt{s_{NN}}$ (in GeV)	2760	3.97 ($E_{lab}\approx 8A$)	8.1 ($E_{lab}\approx 35A$)
$\langle n_{ch}\rangle^{min.bias}_{our\ model}$	8895	328.75	670.8
$\langle n_{ch}\rangle^{central}_{our\ model}$	14377	380.8	837
$\langle n_{ch}\rangle^{central}$ [6,79]	≈ 13124 ($N_{part}=386$)	–	–
$\langle n_{\pi}\rangle^{central}_{HSD}$ [77,78]	–	≈ 308	≈ 650

In Fig. 5, we plot the variations of $(dn_{ch}/d\eta)_{\eta=0}$ obtained in our model for $Au-Au$, $Pb-Pb$ and $Cu-Cu$ central collisions with respect to $\sqrt{s_{NN}}$. We also compare our model results with the experimental data of RHIC, SPS and LHC experiments [62,63,64,65,66,73,74]. Our model results compare well with the experimental data from very low energy to very high energy. As we didn't take any effect of final-state interactions in our model, it hints at the fact that the hadron multiplicity in $A-B$ collisions is mainly driven by initial parton production and the effect of final-state interaction should be negligibly small as suggested in some earlier calculations [75,76].

In Table II, we give the predictions for the mean multiplicity of charged hadrons, produced in minimum-bias events as well as in central events, based on our model calculations at LHC and CBM energies. Specifically, we calculate the charged hadron multiplicity in $Pb-Pb$ collisions at $\sqrt{s_{NN}}=2.76$ TeV for LHC and the mean multiplicity of charged hadrons in $Au-Au$ collisions at $E_{lab} = 8$ AGeV and 35 AGeV for future CBM experiment. We also compare our model results with the other model predictions like hadron string dynamics (HSD) model for CBM experiment [77,78] and model of W. Busza for LHC experiment [6,79], respectively.

4 Conclusions and Future Prospects

Within the framework of constituent quarks, we have given a parametrization which interrelates the multiplicity and mid-rapidity density distributions in p-p, p-A and A-A collisions from a few GeV upto the highest LHC energies. It involves the interactions of the constituent quarks of the colliding objects or beam and target particle or nuclei and the mean number of collisions suffered by them. We assume that the production of secondary particles is proportional to the fraction of the available energy for the participant quarks. This picture describes consistently the 'soft' hadron production in $p-p$, $p-\bar{p}$, $p-A$ and $A-A$ collisions. In heavy-ion collisions, one participating quark can have larger number of interactions due to the large size of the nucleus as well as due to a large travel path to be travelled inside the nucleus and hence, we get larger energy available for the secondary particle production [29]. We find that our parametrization gives an excellent fit to the $p-p$ data in the entire energy range. We also find that a suitable extension of this description works very well for the nucleus-nucleus $(A-A)$ collisions in the entire energy range. We have also compared our model results with the results obtained from modified Glauber model to demonstrate the difference between the role played by nucleon-nucleon interactions vis-a-vis quark-quark interactions. We have also shown the charged particle multiplicity in central as well as in minimum-bias $d-Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV and we find the results compare quite well with the experimental results. This exercise clearly shows that our model also works quite well for asymmetric colliding nuclei case. We have also proposed an extension of the two-component model based on our parameterization and the results for rapidity distribution obtained by us again work well. We have given our predictions for LHC and CBM data as well and compare them with other model predictions [6,79] existing in the literature.

In conclusion, we have attempted to point out a mechanism for 'soft' hadron production common to hadron-hadron, hadron-nucleus and nucleus-nucleus collisions. Due to non perturbative nature, QCD is not expected to work in these cases. So lacking a workable theory, we have relied on the phenomenology and all the salient features of the experimental data are very well explained by our parameterization. We hope that the work done here will highlight the multiparticle production mechanism present in nucleus-nucleus collisions in relation to p-p interactions. This can help in detecting any deviation observed in the data from the predictions of our model and thus a hint for QGP formation would appear likely in undergoing ultra relativistic heavy-ion experiments. Furthemore, future extension of this scheme can help us to study the rapidity, pseudorapidity as well as transverse mass distributions of the charged hadrons in the entire rapidity region and this can form a matter for future publication.

Acknowledgments

PKS is grateful to the University Grants Commission (UGC), New Delhi for providing a research grant. BKS also sincerely acknowledges the Indian Space Research Organization (ISRO) and Department of Science & Technology (DST), India for providing the financial support.

References

1. C. P. Singh, Phys. Rep. 236 (1993) 147.
2. W. Kittel, E. A. De Wolf, Soft Multihadron Dynamics (World Scientific, 2005), 652 pp.
3. J. Manjavidze, A. Sissakian, Phys. Rep. 346 (2001) 1; I. M. Dremin, J. W. Gary, Phys. Rep. 349 (2001) 301.
4. N. S. Amelin, N. Armesto, C. Pajares, D. Sousa, Eur. Phys. J. C 22 (2001) 149.
5. V. Topor Pop, M. Gyulassy, J. Barrette, C. Gale, S. Jeon and R. Bellwied, Phys. Rev. C 75 (2007) 014904.
6. N. Armesto et al., J. Phys. G 35 (2008) 054001.
7. H. J. Drescher, M. Hladik, S. Ostapchanko, T. Pierog and K. Werner, Phys. Rep. 350 (2001) 93.
8. Z. W. Lin, C. M. Ko, B. A. Li, B. Zhang and S. Pal, Phys. Rev. C 72 (2005) 064901.
9. X. N. Wang, M. Gyulassy, Phys. Rev. D 44 (1991) 3501.
10. I. P. Lokthin, L. V. Malinina, S. V. Petrushanko, A. M. Snigirev, I. Arsene and K. Tywoniuk, arXiv:0903.0525 [hep-ph] (2009).
11. M. Mitrovski, T. Shuster, G. Graef, H. Petersen and M. Bleicher, Phys. Rev. C 79 (2009) 044901.
12. T. Sjostrand, S. Mrenna, P. Z. Skands, JHEP 0605 (2006) 026.
13. F. W. Bopp, J. Ranft, R. Engel, S. Roesler, Phys. Rev. C 77 (2008) 014904.
14. B. H. Sa, D. M. Zhou, B. G. Dong, Y. L. Yan, H. L. Ma and X. M. Li, J. Phys. G 36 (2009) 025007 .
15. J. Dias de Deus, J. G. Milhano, Nucl. Phys. A 795 (2007) 98.
16. J. L. Albacete, Phys. Rev. lett. 99 (2007) 262301.
17. N. Armesto, C. A. Salgedo, U. A. Wiedemann, Phys. Rev. Lett 94 (2005) 022002.
18. D. Kharzeev, E. Levin, M. Nardi, Nucl. Phys. A 747 (2005) 609.
19. R. Kuiper, G. Wolschin, Annalen Phys. 16 (2007) 67.
20. A. K. Choudhary, Phys. Lett. B 672 (2009) 126.
21. A. Capella, A. Kaidalov, J. Tran Thanh Van, Heavy Ion Phys. 9 (1999) 169.
22. A. El, Z. Xu, C. Greiner, Nucl. Phys. A 806 (2008) 287.
23. A. Bzdak, Acta Phys. Polon. B 39 (2008) 1977.
24. A. Bialas, M. Bleszynski, W. Czyz, Nucl. Phys. B 111 (1976) 461.
25. T. J. Humanic, Phys. Rev. C 79 (2009) 044902.
26. E. K. G. Sarkisyan, A. S. Sakharov, AIP Conf. Proc. 828 (2006) 35.

27. K. Aamodt et al. (ALICE Collaboration), Phys. Rev. Lett 105 (2010) 252301.
28. C. P. Singh, M. Shyam, S. K. Tuli, Phys. Rev. C 40 (1989) 1716; M. Shyam, C. P. Singh and S. K, Tuli, Phys. Lett. B 164 (1985) 189; C. P. Singh, and M. Shyam, Phys. Lett. B 171 (1986) 125; C. P. Singh, M. Shyam and S. K. Tuli, ICTP Report No. IC/86/159 (1986).
29. E. K. G. Sarkisyan, A. S. Sakharov, Eur. Phys. J. C 70 (2010) 533.
30. E. Fermi, Prog. Theor. Phys. 5 (1950) 570.
31. C.-Y. Wong, Phys. Rev. C 78 (2008) 054902.
32. L. D. Landau, Izv. Akad. Nauk. SSSR 17 (1953) 51.
33. E. Albini et. al., Nuovo Cimento A 32 (1976) 101.
34. A. Breakstone et al., Phys. Rev. D 30 (1984) 528.
35. G. Arnison et. al. Phys. Lett. B 123 (1983) 108; C. Albajar et. al. Nucl. Phys. B 335 (1990) 261.
36. R. E. Ansorge et. al., Z. Phys. C 43 (1989) 357.
37. G. J. Alner et. al., Phys. Lett. B 160 (1985) 193; Phys. Rep. 154 (1987) 247.
38. F. Abe et. al., Phys. Rev. D 41 (1990) 2330.
39. J. Whitmore, Phys. Rep. 10 (1974) 273.
40. J. F. Grosse-Oetringhaus and K. Reygers, J. Phys. G 37 (2010) 083001.
41. P. Slattery, Phys. Rev. Lett. 29 (1972) 1624.
42. W. Thome et. al., Nucl. Phys. B 129 (1977) 365.
43. S. Jeon, V. T. Pop and M. Bleicher, Phys. Rev. C 69 (2004) 044904.
44. A. Z. Patashinskii, JETP Lett. 19 (1974) 338.
45. M. Ta-chung, Phys. Rev. D 15 (1977) 197.
46. V. V. Anisovich and V. M. Shekhter, Nucl. Phys. B 55 (1973) 455; V. V. Anisovich et al., Nucl. Phys. B 133 (1978) 477.
47. A. Bialas et al., Acta Phys. Pol. B 8 (1977) 585.
48. G. V. Devidenko, N. N. Nikoloev, Sov. J. Nucl. Phys. 24 (1976) 402.
49. V. V. Anisovich, N. M. Kobrinsky, J. Nyiri, Yu. M. Shabelsky, Quark Model and High Energy Collisions (World scientific, Singapore, 2004).
50. J. Nyiri, Int. J. Mod. Phys. A 18 (2003) 2403.
51. M. Shyam, S. K. Tuli, Z. Phys. C 29 (1985) 267.
52. M. L. Miller et. al., Annu. Rev. Nucl. Part. Sci. 57 (2007) 205.
53. H. DeVries, C. W. DeJager, and C. DeVries, Atomic Data and Nuclear Data Tables 36 (1987) 495.
54. M. Basile et al., Nuovo Cimento A 66 (1981) 129.
55. M. Basile et al., Nuovo Cimento A 73 (1983) 329.
56. T. F. Huang et al., Z. Phys. C 29 (1985) 611.
57. X. N. Wang, and M. Gyulassy, Phys. Rev. Lett. 86 (2001) 3496.
58. A. Capella and D. Sousa, Phys. Lett. B 511 (2001) 185.
59. K. J. Eskola, K. Kajantie, P. V. Ruuskanen, and K. Tuominen, Nucl. Phys. B 570 (2000) 379.
60. D. Kharzeev and M. Nardi, Phys. Lett. B 507 (2001) 121.
61. G. Feofilov and A. Ivanov, Journal of Physics: Conference Series 5 (2005) 230.
62. B. Alver et. al., Phys. Rev. C 83 (2011) 024913.
63. S. S. Adler et. al., Phys. Rev. C 71 (2005) 034908.
64. B. B. Back et. al., Phys. Rev. C 74 (2006) 021902 (R).
65. J. L. Klay et. al., Phys. Rev. C 68 (2003) 054905.
66. S. V. Afanasiev et. al., Phys. Rev. C 66 (2002) 054902.
67. B. B. Back et. al. (PHOBOS Coll.), Phys. Rev. C 72 (2005) 031901.
68. J.-X. Sun, F.-H. Liu, E.-Q. Wang, Y. Sun, and Z. Sun, Phys. Rev. C 83 (2011) 014001.
69. R. Nouicer et. al., J. Phys. G 30 (2004) S1133.
70. B. I. Abelev et. al., Phys. Rev. C 79 (2009) 034909.
71. K. Aamodt et. al., Eur. Phys. J. C 68 (2010) 89 ; Eur. Phys. J. C 65 (2010) 111.
72. V. Khachatryan et. al., J. High Energy Phys. 2010 (2010) 02041; Phys. Rev. Lett. 105 (2010) 022002.
73. K. Aamodt et. al., Phys. Rev. Lett. 106 (2011) 032301.
74. S. Chatrchyan et. al., J. High Energy Phys. 08 (2011) 141.
75. K. J. Eskola, V. J. Kolhinen, and C. A. Salgado, Eur. Phys. J. C 9 (1999) 61.
76. Z. W. Lin, S. Pal, C. M. Ko, B. A. Li, and B. Zhang, Phys. Rev. C 64 (2001) 011902.
77. CBM Report 2012-01, Nuclear Matter Physics at SIS-100, http://www.gsi.de/documents/Doc-2011-Aug-29-1.pdf.
78. W. Cassing, E. L. Bratkovskaya, S. Juchem, Nucl. Phys. A 674 (2000) 249.
79. W. Busza, Acta. Phys. Polo. B 35 (2004) 2873.